## 4.6. Prevalence of Selected Chronic Conditions By Age and Gender - Per 1,000 Persons 1,2 in Hawai`i, Hawai`i Health Survey (HHS) 1998

Demographic _ Characteristic	Age in Years									Total <sup>3</sup>	
	0-14	15-17	18-24	25-34	35-44	45-54	55-64	65-74	75+	N	Prevalence/ per 1,000 Persons
Condition Type					Female						
Arthritis	0.7*	7.2*	9.9	18.4	41.5	123.4	224.9	340.3	393.2	56,745	97.9
Asthma	114.3	79.2	85.5	72.8	86.7	85.3	69.0	73.0	66.6	50,149	86.5
Diabetes	2.7*	1.8*	13.8	15.4	36.0	69.3	131.1	121.9	134.4	28,059	48.4
High Blood Cholesterol	3.5*	14.3*	15.3	42.2	87.5	178.0	316.9	393.6	300.1	71,969	124.1
Hypertension	4.2	1.8*	22.2	45.9	77.0	187.9	301.7	390.9	393.4	74,442	128.4
Total	125,579	22,942	48,752	74,824	98,540	77,550	49,131	48,281	34,297	579,896	
Condition Type					Male						
Arthritis			5.2	11.2	28.6	62.9	117.7	164.2	225.2	27,556	48.4
Asthma	167.1	147.4	114.0	57.7	58.1	69.1	70.6	32.5	66.1	52,384	92.1
Diabetes	0.9*		3.2*	9.9	21.1	85.5	122.8	151.1	116.9	24,937	43.8
High Blood Cholesterol	4.3	2.0*	17.0	65.1	112.4	233.8	320.8	362.8	239.3	72,058	126.7
Hypertension	1.5*	10.6*	11.8	63.8	115.5	235.8	313.0	370.4	377.0	76,023	133.6
Total	128,268	20,515	52,232	76,613	97,936	77,269	45,588	40,576	29,917	568,914	1,148,810

<sup>&</sup>lt;sup>1</sup> All household members

Sample size 12,923 people

<sup>&</sup>lt;sup>2</sup> Sample numbers provisionally weighted and adjusted\* for total population of Hawai`i

<sup>\*(</sup>total numbers are adjusted as Ni`ihau, Kalawao, group quarters, and homeless are not represented, see HHS Procedure Manual 1998)

<sup>&</sup>lt;sup>3</sup> Row and/or column totals may not sum to totals listed due to rounding

Number of cases in sample numerator or denominator too small for reliability - shown for consistency only

<sup>--</sup> No reported cases